



Supplementary Fig. 1. CONSORT

Supplementary Table 1. Comparison of different degrees of renal injury with various clinical indicators

Indicators	Mild	Moderate	Severe	<i>P</i> values
FENDRR	1.29 ±0.15	1.62 ±0.06	1.89 ±0.33	< 0.001
PCT (ng/ml)	4.38 ±2.99	6.48 ±3.14	6.81 ±4.52	0.013
BUN (mmol/l)	28.09 ±19.14	36.76 ±12.72	40.87 ±22.78	0.030
SCr (μmol/l)	297.44 ±171.43	371.36 ±207.88	489.65 ±168.94	0.001
APACHE II	16.38 ±1.06	17.19 ±7.20	21.68 ±5.76	0.003
SOFA	9.23 ±4.39	11.08 ±3.27	12.00 ±3.61	0.031
Lactate (mmol/l)	3.61 ±1.98	4.59 ±1.70	5.04 ±2.23	0.020
Length of hospital stay (days)	16.13 ±5.71	18.00 ±6.63	19.16 ±6.57	0.183

PCT – procalcitonin, BUN – blood urea nitrogen, SCr – serum creatinine, APACHE II – Acute Physiology and Chronic Health Evaluation II, SOFA – Sequential Organ Failure Assessment